

Relativistic corrections to the Ekert test for eavesdropping

Marek Czachor

*Katedra Fizyki Teoretycznej i Metod Matematycznych
Politechnika Gdańska, ul. Narutowicza 11/12, 80-952 Gdańsk, Poland*

A degree of violation of the Bell inequality depends on momenta of massive particles with respect to a laboratory if spin plays a role as a “yes–no” observable. For ultra-relativistic particles a standard Ekert test has to take into account this velocity dependent suppression of the degree of violation of the inequality. Otherwise Alice and Bob may “discover” a nonexistent eavesdropper.

The cryptographic protocol proposed by Ekert [1] consists of two essential parts. First, a two-particle singlet-state source produces pairs of particles generating identical sequences of bits which can be subsequently used by Alice and Bob as a cryptographic key. Second, some randomly chosen data are used to test whether the particles were really prepared in a singlet state. This can be done by measuring the “Bell average”

$$c(a, a', b, b', \psi) = \langle \psi | \hat{a} \otimes \hat{b} + \hat{a} \otimes \hat{b}' + \hat{a}' \otimes \hat{b} - \hat{a}' \otimes \hat{b}' | \psi \rangle$$

where the binary observables $\hat{a}$, etc., are chosen in such a way that the Bell inequality $|c(a, a', b, b', \psi)| \leq 2$ is violated. This will be done most efficiently if $|c(a, a', b, b', \psi)| = 2\sqrt{2}$ and the violation is expected to be maximal. A result of the form $|c(a, a', b, b', \psi)| < 2\sqrt{2}$ means that $|\psi\rangle$ is not a pure singlet, and this indicates an eavesdropper. After having detected the eavesdropper the data collected by Alice and Bob are thrown away and the whole procedure starts again. The binary observables appearing in $c(\dots)$ represent results of some “yes–no” experiments and it is known

that for any entangled state there exists a set of observables leading to the maximal violation of the inequality. This statement is, strictly speaking, true *in principle*. In practice, however, there is no operational rule that could tell us how to measure an arbitrary observable. So practically we are confined to a concrete set of “physical” observables which, in this context, are various “polarizations”. In the photon case the “polarizations” are either simply linear polarizations or some observables related to interferometric experiments. The situation is less clear if one considers massive particles. At this moment one may ask why should one worry about massive particles if massless photons work

so well. A practical answer is provided by a very high efficiency of atomic detectors as compared to the photonic ones [2]. Taking into account the spectacular comeback of atoms in interferometry and the recent progress in experiments with coherent atomic sources one cannot exclude future applications of atomic entangled states in cryptography. The question I want to discuss here is the role of relativistic effects that may become important if very fast atomic EPR-type beams are used for the cryptographic key distribution [3]. Let us begin with the yes–no polarizations for

relativistic massive particles. It is known that the spinor part $\vec{s}$ of the generator of rotations is not a well defined spin operator for, say, the Dirac electrons. Its projection on the momentum direction (helicity) commutes with the free Dirac Hamiltonian, but to measure $c(\dots)$ we need measurements of spin in different directions. The naive approach based on $\hat{a} = \vec{a} \cdot \vec{s}$ cannot be physically meaningful since eigenstates of $\hat{a}$ are not preserved by a free evolution. This is a general phenomenon present in all Poincaré covariant systems but this argument alone does not prove that any notion of spin is meaningless. There are at least two objects whose definitions are representation-independent and which can represent a physical spin of a relativistic system. One is the spin operator $\vec{S}$ constructed by means of a relativistic center-of-mass position operator. The other is the Pauli-Lubanski vector $(W_0, \vec{W})$. The two objects are not equivalent but their spacelike parts are proportional to each other. Their projections on $\vec{a}$ have different eigenvalues but the same eigenstates. Accordingly they both give the same values of $c(\dots)$. Let $a_\alpha = (a_0, \vec{a})$ be some

world-vector. One can consider joint eigenstates of the Pauli-Lubanski vector and the 4-momentum. The eigenvalues of $a_\alpha W^\alpha = a \cdot W$ are

$$w_\pm(a, p) = \pm j_3 \sqrt{(a \cdot p)^2 - a^2 p^2} = \pm w(a, p)$$

where $j_3 = 1/2, 1, 3/2, \dots, j$, and $a^2 = a_\alpha a^\alpha$ etc. For $j = 1/2$ the “yes–no” observable $\hat{a}$ can be defined in a relativistically invariant way as

$$\hat{a} = \hat{a}(p) = \frac{a \cdot W}{w(a, p)}.$$

An arbitrary two-spin-1/2-particle state can be written as

$$|f\rangle = \sum_{\sigma_1, \sigma_2} \int d\Gamma(\vec{p}_1) d\Gamma(\vec{p}_2) f(\sigma_1, \vec{p}_1, \sigma_2, \vec{p}_2) |\sigma_1, \vec{p}_1\rangle |\sigma_2, \vec{p}_2\rangle$$

where σ_k is a helicity, $f(\sigma_1, \vec{p}_1, \sigma_2, \vec{p}_2) = -f(\sigma_2, \vec{p}_2, \sigma_1, \vec{p}_1)$, and $d\Gamma(\vec{p}) = (2\pi)^{-3} (2|p_0|)^{-1} d^3p$ is an invariant measure on a mass hyperboloid. Let us consider the particular state

$$|f\rangle = \frac{1}{\sqrt{2}} \int d\Gamma(\vec{p}_1) d\Gamma(\vec{p}_2) f(\vec{p}_1, \vec{p}_2) \left(|+, \vec{p}_1\rangle |-, \vec{p}_2\rangle - |-, \vec{p}_1\rangle |+, \vec{p}_2\rangle \right) \quad (1)$$

where $f(\vec{p}_1, \vec{p}_2) = f(\vec{p}_2, \vec{p}_1)$. Denote $\vec{\beta} = \vec{p}/p_0$, $\vec{n} = \vec{p}/|\vec{p}|$, and consider projections of W^α on spacelike directions satisfying in a laboratory frame $a_\alpha = (0, \vec{a})$, $b_\alpha = (0, \vec{b})$. The laboratory frame is here the one which defines the decomposition of the 4-momentum into energy and 3-momentum $\vec{p}$ appearing in the wave functions. The EPR average is

$$\begin{aligned} \langle f | \hat{a} \otimes \hat{b} | f \rangle &= - \int d\Gamma(\vec{p}_1) d\Gamma(\vec{p}_2) |f(\vec{p}_1, \vec{p}_2)|^2 \\ &\times \frac{\left(\sqrt{1 - \beta_1^2} [\vec{a} - (\vec{a} \cdot \vec{n}_1) \vec{n}_1] + (\vec{a} \cdot \vec{n}_1) \vec{n}_1 \right) \cdot \left(\sqrt{1 - \beta_2^2} [\vec{b} - (\vec{b} \cdot \vec{n}_2) \vec{n}_2] + (\vec{b} \cdot \vec{n}_2) \vec{n}_2 \right)}{\sqrt{1 + \beta_1^2 [(\vec{n}_1 \cdot \vec{a})^2 - 1]} \sqrt{1 + \beta_2^2 [(\vec{n}_2 \cdot \vec{b})^2 - 1]}} \end{aligned} \quad (1)$$

For

$$d\Gamma(\vec{p}_1) d\Gamma(\vec{p}_2) |f(\vec{p}_1, \vec{p}_2)|^2 \approx d^3p_1 d^3p_2 \delta(\vec{p} - \vec{p}_1) \delta(\vec{p}_1 - \vec{p}_2) \quad (2)$$

we obtain the result discussed in [3]:

$$\langle f | \hat{a} \otimes \hat{b} | f \rangle \approx - \frac{\vec{a} \cdot \vec{b} - \beta^2 \vec{a}_\perp \cdot \vec{b}_\perp}{\sqrt{1 + \beta^2 [(\vec{n} \cdot \vec{a})^2 - 1]} \sqrt{1 + \beta^2 [(\vec{n} \cdot \vec{b})^2 - 1]}}. \quad (3)$$

(3) implies

$$\langle f | \hat{a} \otimes \hat{a} | f \rangle = -1,$$

which means that spins are always anti-parallel. The state (1) satisfying (2) can be therefore used for a secure key transfer. The same is true for more realistic states satisfying

$$d\Gamma(\vec{p}_1) d\Gamma(\vec{p}_2) |f(\vec{p}_1, \vec{p}_2)|^2 \approx d^3p_1 d^3p_2 |f(\vec{p}_1)|^2 \delta(\vec{p}_1 - \vec{p}_2)$$

where $|f(\vec{p}_1)|^2$ is a probability density. Notice that for $\vec{a} = \vec{a}_\perp$, $\vec{b} = \vec{b}_\perp$, we obtain the nonrelativistic formula

$$\langle f | \hat{a} \otimes \hat{b} | f \rangle = -\vec{a} \cdot \vec{b}.$$

The condition $\vec{a} = \vec{a}_\perp$, etc., cannot be satisfied in a real experiment since $\vec{a}$ is a fixed vector and the set of momenta perpendicular to a given fixed axis has 3-dimensional measure 0. So from any realistic wave packet one can remove all $\vec{p}$ perpendicular to $\vec{a}$ and an experimentally measured average will not change. So assume that $\vec{a} \cdot \vec{p} \neq 0$. In this case we find that

$$\langle f | \hat{a} \otimes \hat{b} | f \rangle \rightarrow -1$$

for $\beta \rightarrow 1$ independently of $\vec{a}$, $\vec{b}$. Here β is the average velocity of the wave packet. To better understand this result

let us recall the explicit form of the relativistic spin operator

$$\vec{S} = \vec{W}/p_0 = \sqrt{1 - \beta^2} \vec{s}_\perp + (\vec{n} \cdot \vec{s}) \vec{n}.$$

In the $\beta \rightarrow 1$ limit the vector points in the momentum direction and

$$[S_k, S_l] \rightarrow 0.$$

The relativistic spin becomes the more “classical” the faster the particle. This explains qualitatively why one should expect a *suppression of the Bell inequality violation* for ultra-relativistic massive particles [3]. It has been shown in [3] that for $\beta \rightarrow 1$

$$|c(a, a', b, b', f)| \rightarrow 2$$

from above even if the measuring devices are optimally chosen.

The plots illustrate these phenomena. The vectors in $c(\dots)$ are chosen coplanarly in a way which leads to the maximal violation of the Bell inequality in the nonrelativistic case ($|c| = 2\sqrt{2}$): $\vec{a} = (1/\sqrt{2}, 1/\sqrt{2}, 0)$, $\vec{a}' = (-1/\sqrt{2}, 1/\sqrt{2}, 0)$, $\vec{b} = (0, 1, 0)$, and $\vec{b}' = (1, 0, 0)$. The measuring devices are at rest in a laboratory frame and $\vec{\beta}$ denotes velocities of particles with respect to the lab.

One may wonder why do we have here a suppression of $|c(a, a', b, b', f)|$ for any spin and simultaneously have maximal violation for photons. The answer is that in the photon polarization or interferometric experiments we do not measure spins. So a possibility exists that there are also some other observables that could be used in the EPR-type experiments with massive particles instead of spin, but none have been proposed so far.

The moral following from our story is that Alice and Bob do not necessarily have to throw away all data if they find $|c(a, a', b, b', f)| < 2\sqrt{2}$. They first have to check the relativistic corrections. In order to do this they have to know the momentum distribution of the particle beam. The momentum can be measured, in principle, on each particle without disturbing its spin since 4-momentum and W_α commute.

ACKNOWLEDGMENTS

This work was partially supported by the KBN grant 2P30B03809 and is a part of the Polish-Flamish project on “New models of probability calculus and new experiments on single quantum particles”. I’m indebted to the Batory Foundation and Nokia-Poland for financial support.




FIG. 1. Both particles move with velocity $\vec{\beta} = \beta(\cos \phi, \sin \phi, 0)$. The maximal violation is found for $\beta = 0$ and there is no violation for $\beta \rightarrow 1$.




FIG. 2. Similar situation, but now the particles move with velocity $\vec{\beta} = \beta(\sin \theta, 0, \cos \theta)$. The maximal violation is found for $\beta = 0$ and is suppressed as β approaches 1. For $\theta = 0$ the particles move perpendicularly to the measuring devices and the violation remains maximal for all β . The set of all such momenta is of measure 0 which implies that one should expect the relativistic damping effect to be present for all wave packets.




FIG. 3. Situation from Fig. 1 but with $\vec{\beta} = \beta(\cos \phi \sin \theta, \sin \phi \sin \theta, \cos \theta)$ for $\beta = 0.99$ (upper) and $\beta = 0.95$ (lower); $0 \leq \phi \leq 2\pi$, $0 \leq \theta \leq \pi$.




FIG. 4. The Bell average with the same choice of measuring devices but now the velocities of the two particles are different: One moves nonrelativistically ($\beta_1 \approx 0$) and the other with an arbitrary velocity $\vec{\beta}_2 = \beta(\cos \phi, \sin \phi, 0)$. The plot is very similar to the one from Fig. 1.




FIG. 5. A cut through the surface from Fig. 1 at the “gully” $\phi = 0$.




FIG. 6. The relativistic corrections are much more visible if instead of measuring the entire Bell average one concentrates on $\langle f|\hat{a} \otimes \hat{b}|f \rangle$ and takes $\vec{a}$ and $\vec{b}$ perpendicular. For particles moving with velocity $\vec{\beta} = \beta\vec{n}$, $\vec{a} \cdot \vec{b} = 0$, $\vec{a} \cdot \vec{n} = \vec{b} \cdot \vec{n} = 2^{-1/2}$ we find $\langle f|\hat{a} \otimes \hat{b}|f \rangle = -\beta^2(2 - \beta^2)^{-1}$. The solid curve shows this average as compared to $[1 - \beta^2]^{1/2} - 1$ (dotted).

-
- [1] A. Ekert, “Quantum cryptography based on Bell’s theorem”, *Phys. Rev. Lett.* **67**, pp. 661-664 (1991).
 - [2] E. S. Fry, T. Walther, and S. Li, “Proposal for a loophole-free test of the Bell inequalities”, *Phys. Rev. A* **52**, pp. 4381-4395 (1995).
 - [3] M. Czachor, “Einstein-Podolsky-Rosen-Bohm experiment with relativistic massive particles”, *Phys. Rev. A* **55**, pp. 72-76 (1997).